

Table 1. Summary of ACIP vaccine recommendations for HCP

Vaccine	Schedule	HCP who should receive vaccination
Hepatitis B (Recombivax HB, Engerix-B, and Twinrix)	Three-dose series: 0, 1, and 6 months	HCP at risk for exposure to blood or body fluids or postexposure prophylaxis
Influenza	One dose annually	All HCP
Tdap	One dose if not previously vaccinated	All HCP should receive a single dose of Tdap
Varicella	Two doses separated by 4 weeks	All HCP must either receive the vaccine or provide proof of immunity which can include: (1) written documentation of two doses of vaccine, (2) laboratory evidence, (3) diagnostic history of varicella or herpes zoster by health care provider.
MMR	Two doses separated by 4 weeks	HCP who lack documentation of two doses of vaccine or laboratory evidence of immunity
Meningococcal (if age <55 years, use MCV4; if ≥55 years, use MPSV4)	Initial two-dose series and revaccination every 5 years	HCP at high risk; not routine for all HCP

Abbreviations used: ACIP, Advisory Committee on Immunization Practices; HCP, health care personnel; MCV4, quadrivalent meningococcal conjugate vaccine; MMR, measles-mumps-rubella; MPSV4, meningococcal polysaccharide vaccine; Tdap, tetanus-diphtheria-acellular pertussis.